AESTHETIC ENHANCEMENTS AT OPFP

XEOMIN® Post Treatment Instructions

1. The full effect of XEOMIN® may take anywhere from 2-14 days.
2. Do not massage, rub or apply pressure to the treated areas for the first 24 hours after treatment. This may cause the XEOMIN® to migrate to other muscles and can cause temporary dropping of the eyelids.
3. Stay erect; do not lie down for at least 4 hours after the injections.
4. Do not exercise for 24 hours and resume activity slowly and carefully after the injections.
5. Bruising may develop at injection site. You may use ice packs at the injection site for bruising and/or discomfort. The ice should be held only in light contact with the skin.
6. Avoid aspirin, ibuprofen, and drinking alcohol for at least 24 hours post treatment.
7. An immediate headache is common especially, if it is your first cosmetic injection treatment.
8. If drooping of eyelids does occur, it is temporary and generally resolves within two to three weeks.
9. Makeup can be reapplied after treatment
10. Increase water intake
11. Seek medical care if problems with the following occur: swallowing, speech, respiratory, or blinking.
12. Avoid driving a vehicle or other hazardous activities if the following occur: loss of muscle strength, blurred vision, or drooping eyelids.

Please call our office with any questions OR if any of the following occur: skin rash, sudden fever, increased pain, increased swelling, or bleeding at any of the injection sites at (716) 508-0160.